

## HEDIS Provider Guide: Cervical Cancer Screening (CCS)

## **Measure Description**

Members 21 to 64 years of age who were recommended for routine cervical cancer screening and were screened for cervical cancer using any of the following criteria:

- Members 21 to 64 years of age who were recommended for routine cervical cancer screening and had cervical cytology performed within the last three years.
- Members 30 to 64 years of age who were recommended for routine cervical cancer screening and had cervical high-risk human papillomavirus (hrHPV) testing performed within the last 5 years.
- Members 30 to 64 years of age who were recommended for routine cervical cancer screening and had cervical cytology/high-risk human papillomavirus (hrHPV) co-testing within the last 5 years.

Members recommended for Cervical Cancer Screening:

- Administrative Gender of Female at any time in the member's history.
- Sex Assigned at Birth of Female at any time in the member's history.
- Sex Parameter for Clinical Use of Female during the measurement period.

**Note:** It is recommended that transgender and gender diverse patients who have a cervix have regular cervical pap tests, as per the published guidelines for cisgender women.

**Exclusions:** Hysterectomy with no residual cervix or cervical agenesis or acquired absence of cervix any time during the member's history through December 31 of the measurement year. Members in hospice or palliative care. Members with Sex Assigned at Birth of Male at any time in the patient's history.

## **Using Correct Billing Codes**

Description	Codes
Codes to Identify Cervical Cancer Screening	
Cervical Cancer Screening	CPT: 88141-88143, 88147, 88148, 88150, 88152, 88153, 88164-88167, 88174, 88175
	HCPCS: G0123, G0124, G0141, G0143-G0145, G0147, G0148, P3000, P3001, Q0091
	<b>LOINC:</b> 10524-7, 18500-9, 19762-4, 19764-0, 19765-7, 19766-5, 19774-9, 33717-0, 47527-7, 47528-5
Codes to Identify HPV Screening	
HPV Screening	<b>CPT</b> : 87624, 87625
	HCPCS: G0476
	<b>LOINC:</b> 21440-3, 30167-1, 38372-9, 59263-4, 59264-2, 59420-0, 69002-4, 71431-1, 75694-0, 77379-6, 77399-4, 77400-0, 82354-2, 82456-5, 82675-0, 95539-3
Exclusions: Hysterectomy with No Residual Cervix	
Cervical Agenesis or Acquired Absence of Cervix Diagnosis	ICD10CM: Q51.5, Z90.710, Z90.712
	ICD9PCS: 752.43, V88.01, V88.03
Hysterectomy with no Residual Cervix	<b>CPT:</b> 57530, 57531, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290–58294, 58548, 58550, 58552–58554, 58570–58573, 58575, 58951, 58953, 58954, 58956, 59135
	ICD10PCS: OUTC0ZZ, OUTC4ZZ, OUTC7ZZ, OUTC8ZZ
	ICD9PCS: 68.41, 68.49, 68.51, 68.59, 68.61, 68.69, 68.71, 68.79, 68.8

## How to Improve HEDIS® Scores

- Request a focused list of members who need a pap test from your Blue Shield Promise Quality Program Manager.
- Use a reminder/recall system.
- Request pap test results be sent to you if done during the OB/GYN visit.
- Document in the medical record if the patient has had a hysterectomy with no residual cervix.
- Do not miss opportunities. For example, complete pap tests during regularly scheduled well woman visits, sick visits, urine pregnancy tests, UTI and Chlamydia/STI screening.

HEDIS® is a registered trademark for the National Committee of Quality Assurance (NCQA)