



Federal Employee Program.

(eculizumab) SOLIRIS
PRIOR APPROVAL REQUEST

Send completed form to:
FAX: 855-895-3504
FOR URGENT FAX: 844-244-0226

Additional information is required to process your claim for prescription drugs. Please complete the patient portion, and have the prescribing physician complete the physician portion and submit this completed form.

Form with two main sections: Patient Information (required) and Provider Information (required). Includes fields for Date, Patient Name, Date of Birth, Sex, Street Address, City, State, Zip, Patient ID, Provider Name, Specialty, NPI, Office Phone, Office Fax, Office Street Address, City, State, Zip, and Physician Signature.

PHYSICIAN COMPLETES

Soliris (eculizumab)

**Check www.fepblue.org/formulary to confirm which medication is part of the patient's benefit

NOTE: Form must be completed in its entirety for processing

Is this request for brand or generic? [] Brand [] Generic

1. Is the prescriber enrolled in the Soliris REMS program? [] Yes [] No

2. What is the patient's diagnosis?

[] Atypical Hemolytic Uremic Syndrome (aHUS)

a. Does the patient have Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS)? [] Yes [] No

b. Will Soliris be used in combination with another Prior Authorization (PA) medication for atypical hemolytic uremic syndrome (e.g., Ultomiris (ravulizumab-cwvz))? [] Yes* [] No

*If YES, specify the medication: _____

c. Has the patient been on Soliris continuously for the last 4 months, excluding samples? Please select answer below:

[] NO - this is INITIATION of therapy, please answer the following questions:

i. Does the patient have a documented baseline value for serum lactate dehydrogenase (LDH)? [] Yes [] No

ii. Has or will the patient be vaccinated against Neisseria meningitidis at least two weeks prior to initiating therapy? [] Yes [] No*

*If NO, is urgent Soliris therapy indicated for this patient (e.g., the risks of delaying treatment with Soliris outweigh the risk of developing a meningococcal infection)? [] Yes [] No

[] YES - this is a PA renewal for CONTINUATION of therapy, please answer the following questions:

i. Has the patient had a decrease in serum lactate dehydrogenase (LDH) from pretreatment baseline? [] Yes [] No

ii. Has the patient experienced unacceptable toxicity while on Soliris therapy? [] Yes [] No

[] Neuromyelitis Optica Spectrum Disorder (NMOSD)

a. Has the patient been on Soliris continuously for the last 4 months, excluding samples? Please select answer below:

[] NO - this is INITIATION of therapy, please answer the following questions:

i. Is the patient anti-aquaporin-4 (AQP4) antibody positive? [] Yes [] No

ii. Has or will the patient be vaccinated against Neisseria meningitidis at least two weeks prior to initiating therapy? [] Yes [] No*

*If NO, is urgent Soliris therapy indicated for this patient (e.g., the risks of delaying treatment with Soliris outweighs the risk of developing a meningococcal infection)? [] Yes [] No

[] YES - this is a PA renewal for CONTINUATION of therapy, please answer the following questions:

i. Has the patient had fewer relapses while on Soliris therapy? [] Yes [] No

ii. Has the patient experienced unacceptable toxicity while on Soliris therapy? [] Yes [] No

PLEASE PROCEED TO PAGE 2 FOR ADDITIONAL DIAGNOSES



Federal Employee Program.

(eculizumab) SOLIRIS
PRIOR APPROVAL REQUEST

Send completed form to:
FAX: 855-895-3504
FOR URGENT FAX: 844-244-0226

Additional information is required to process your claim for prescription drugs. Please complete the patient portion, and have the prescribing physician complete the physician portion and submit this completed form.

PAGE 2 - PHYSICIAN COMPLETES

Patient Name: _____ DOB: _____ Patient ID: R _____

Generalized Myasthenia Gravis (gMG)

a. Will Soliris be used in combination with another Prior Authorization (PA) C5 complement inhibitor for generalized myasthenia gravis (e.g., Ultomiris (ravulizumab-cwvz))? Yes No

If YES, specify the medication: _____

b. Has the patient been on Soliris continuously for the last 4 months, excluding samples? Please select answer below:

NO - this is INITIATION of therapy, please answer the following questions:

i. Does the patient have a positive serologic test for anti-AChR antibodies? Yes No

ii. What is the patient's MGFA (Myasthenia Gravis Foundation of America) clinical classification? Select answer below:

Class I Class II to IV Class V Unknown

iii. If Class II to IV, does the patient have a documented baseline *MG-Activities of Daily Living (MG-ADL) total score greater than or equal to 6? Yes No

*MG-ADL: http://c.peerview.com/inReview/programs/150204324/downloads/PVI_practiceaids_RMU.pdf

iv. Has or will the patient be vaccinated against Neisseria meningitidis at least two weeks prior to initiating therapy? Yes No*

If NO, is urgent Soliris therapy indicated for this patient (e.g., the risks of delaying treatment with Soliris outweighs the risk of developing a meningococcal infection)? Yes No

v. Does the patient have an intolerance or contraindication or have they had an inadequate treatment response to an acetylcholinesterase inhibitor? Yes No

vi. Does the patient have an intolerance or contraindication or have they had an inadequate treatment response to at least one immunosuppressive therapy either in combination or as monotherapy, such as: azathioprine, cyclosporine, mycophenolate mofetil, tacrolimus, methotrexate, or cyclophosphamide? Yes No

YES - this is a PA renewal for CONTINUATION of therapy, please answer the following questions:

i. Is there a documented decrease of the *MG-Activities of Daily Living (MG-ADL) total score from baseline of greater than or equal to 2 points? Yes No

*MG-ADL: http://c.peerview.com/inReview/programs/150204324/downloads/PVI_practiceaids_RMU.pdf

ii. Has the patient experienced unacceptable toxicity while on Soliris therapy? Yes No

Paroxysmal Nocturnal Hemoglobinuria (PHN)

a. Will Soliris be used in combination with another Prior Authorization (PA) medication for paroxysmal nocturnal hemoglobinuria (e.g., Empaveli (pegcetacoplan), Ultomiris (ravulizumab-cwvz))? Yes No

If YES, specify the medication: _____

b. Has the patient been on Soliris continuously for the last 4 months, excluding samples? Please select answer below:

NO - this is INITIATION of therapy, please answer the following questions:

i. Does the patient have a documented baseline value for serum lactate dehydrogenase (LDH)? Yes No

ii. Has or will the patient be vaccinated against Neisseria meningitidis at least two weeks prior to initiating therapy? Yes No*

If NO, is urgent Soliris therapy indicated for this patient (e.g., the risks of delaying treatment with Soliris outweigh the risk of developing a meningococcal infection)? Yes No

YES - this is a PA renewal for CONTINUATION of therapy, please answer the following questions:

i. Has the patient had a decrease in serum lactate dehydrogenase (LDH) from pretreatment baseline? Yes No

ii. Has the patient experienced unacceptable toxicity while on Soliris therapy? Yes No

None of the above